

Heavy Flavor Physics with Fast MAPS Detector at sPHENIX

Xin Dong

Lawrence Berkeley National Laboratory





Mar. 8, 2017


sPHENIX MVTX Physics Discussion

X. Dong

Uniqueness of Heavy Quarks in QCD




Zhu et al., PLB 647(2007)366




When $M_{HQ} \gg T$, $M_{HQ} \gg gT$

"Brownian" motion $\frac{dp^i}{dt} = -\eta_D p^i + \xi^i(t)$
 → Langevin simu.




Physics Goals of Heavy Flavor Measurements

- Mass/flavor dependence of parton energy loss
- Quantify the medium transport parameter – heavy quark diffusion coefficient, DHQ




PRL 108 (2012) 022301



QCD white paper - arXiv: 1502.02730



Achievements in Charm-Hadron R_{AA}



$R_{AA}(D) \sim R_{AA}(h)$ at $p_T > 4$ GeV/c

- Significant suppression of charmed hadron R_{AA} in central A+A collisions
- strong charm-medium interactions

Achievements in Charm-Hadron v_2



- v_2 of D^0 follows the same trend as light hadrons
- Charm quarks flow the same as light quarks
- Indication of charm quark thermalization

Go Heavier - Open Bottom Production

Open bottom production over a wide range of momentum


Mass/Flavor dependence of parton energy loss

Cleanest probe to quantify medium transport properties – e.g. D_{HQ}

Total bottom yield for precision interpretation of Upsilon suppression

- **low p_T coverage is critical**

Das et al., PRC 90 (2014) 044901



Is charm heavy enough?


Sizable correction to the Langevin approach for charm

- may limit the precision in determining D_{HQ}


Recent Achievements on Bottom

LHC 2.76 TeV

$$R_{AA}(J/\psi_B) > R_{AA}(D)$$





$$R_{AA}(e_B) > R_{AA}(e_{D+B})$$



RHIC 200 GeV

$$R_{AA}(e_B) > R_{AA}(e_D)$$




- R_{AA} of B-meson, b-jet, non-prompt J/ψ 's are consistent with light hadrons/jets at $p_T > 10 \text{ GeV}/c$
- Less R_{AA} suppression for B-decay daughters (J/ψ 's, e) compared to D-mesons at low p_T

Evidence of less energy loss of bottom in the QGP

Requirements for Precision Open Bottom Production at RHIC


- High luminosity runs and large datasets (triggered and untriggered)
 $B \rightarrow J/\psi$, $B \rightarrow D$, $B \rightarrow e$, $B \rightarrow D\pi$ and b -jet etc.
→ **Fast DAQ rate**
- Fast silicon detector with ultimate pointing resolution
Next generation MAPS sensors with much shorter integration time $< 20 \mu\text{s}$ (vs. $186 \mu\text{s}$)
- high efficiency at high RHIC luminosity, particularly at low p_T



Physics Channels

Hadron	Abundance	$c\tau$ (μm)
D^0	56%	123
D^+	24%	312
D_s	10%	150
Λ_c	10%	60
B^+	40%	491
B^0	40%	455
B_s	10%	453
Λ_b	10%	435



$B \rightarrow J/\psi + X$ 1.2%
 $B \rightarrow \bar{D}^0 + X$ 60%
 $B \rightarrow e + X$ 11%
 $B^+ \rightarrow \bar{D}^0 \pi^+$ 0.5% } Needed for
 p_T < 10 GeV
 b-tagged jet



Theory curves on B/D-mesons from TAMU/DUKE/ CUJET



Estimation for Non-Prompt D⁰

- Single track efficiency, DCA distributions taken from full Hijing + GEANT simulations
- These fed into a fast Monte Carlo package to generate the distributions for signals (prompt and non-prompt D⁰) as well as combinatorial background



Physics Performance via Non-Prompt D⁰

100B MB Au+Au collisions at 200 GeV



Assuming signal scales with N_{bin} , background scales with N_{part}^2


To-do list:

- Realistic estimation on the pileup MB/UPC hit density at MAPS sensors.
- Decay channels B->D, B->J/ ψ , B->e and B->D π etc.
- Tracking improvement

Λ_c and HQ Correlations

High statistics Λ_c measurements

Λ_c/D^0 enhancement sensitive to
 - charm quark hadronization,
 thermalization, domains in sQGP etc.




Lee et al, PRL 100 (2008) 222301

Ghosh et al, PRD 90 (2014) 054018

STAR, QM17

Heavy quark correlations



- More sensitivity to HQ-medium interaction, thus better determination of ΔE mechanisms and D_{HQ}
- LHC vs. RHIC – different initial pair correlation/medium dynamics



Zhu et al, PRL 100 (2008) 152301


Backups

D_s Enhancement in Heavy Ion Collisions



- Significant D_s / D^0 enhancement in mid-central Au+Au and Pb+Pb collisions w.r.t fragmentation baseline or p+p measurement
 - Coalescence hadronization
 - SHM predicts D_s / D^0 ratio $\sim 0.35\text{-}0.40$ (central) *A. Andronic et al., PLB 571 (2003) 36*
 - relation to charm quark thermalization in QGP?

Λ_c Enhancement in Heavy Ion Collisions





Ko model : Y. Oh, et.al. PRC 79 (2009) 044905; Greco model : S.Ghosh, et. al. PRD 90 (2014) 054018

- Significant enhancement in Λ_c/D compared to PYTHIA/fragmentation baseline
- The Λ_c/D^0 ratio is compatible with light flavor baryon-to-meson ratios
- Consistent with coalescence + thermalized charm quarks

Summary



- Heavy-flavor phase-I program at RHIC (2014-2016)
 - Precision charmed hadron measurements from STAR-HFT/PHENIX-(F)VTX
- Heavy-flavor phase-II program at RHIC (2021+)
 - Open bottom / correlation measurements
 - Complementary to the HF program at LHC
- Fast MAPS silicon detector is necessary and will deliver the physics



Buzzatti et al., PRL 108 (2012) 022301

arXiv: 1502.02730, 1506.03981

STAR Heavy Flavor Tracker



STAR HFT/PXL – first application of MAPS pixel detector at a collider

- Aim for precision measurements of charmed hadron production in HIC
- PXL detector designed, developed and constructed (including mechanics) at LBNL
- First layer thickness: $0.4\%X_0$
- Pitch size $20.7 \times 20.7 \mu\text{m}$
- Integration time: 186 μs (see next page)


Hit Density on STAR PXL at RHIC Environment

Simulation@50kHz	PXL inner	PXL outer
Radius (cm)	2.8	8
MB pileup hits (cm^{-2})	13	~ 3
UPC electrons (cm^{-2})	33	~ 3
Total bkgd hits (cm^{-2})	46	~ 6
MB signal Au+Au (cm^{-2})	~ 8	~ 1
Au+Au MB real data (cm^{-2})	~ 50	~ 5

Signal hits fraction in MB (Central) events:
 ~15% (~30%) at PXL inner

Increasing fake matches in low p_T



Technology chosen considering both
 physics and technology readiness



Uniqueness at RHIC



Uniqueness at RHIC

- dominated by pair creation, clean interpretation for experimental results



T. Sjostrand, EPJC17 (2000) 137

Fast MAPS Detectors at RHIC – STAR HFT+




STAR HFT+ upgrade / sPHENIX pixel detector:


- Faster ($<20\mu\text{s}$) MAPS sensors – benefiting from ALICE ITS upgrade
- Aim for precision bottom measurements in 2021+ at RHIC

Complementary to LHC heavy flavor program

Pointing Resolution and Efficiency



Single Track Efficiency for the HFT (D0 Efficiency dashed) .vs. Pt



80×10^{26} RHIC luminosity, central Au+Au 200 GeV collisions


MAPS: 20 us integration time

MB pileup events and UPC hits included (small contribution)

Efficiency or good hit probability: inefficiency due to mis-association

$$\text{Efficiency/Good Hit Probability} = \frac{1}{1 + 2\pi R_{r\phi} R_z \rho} \quad R = \sqrt{R_{proj}^2 + R_{det}^2}$$

Tracking Configuration Study



Tracking Configuration Study

R- ϕ Pointing Resolution .vs. Pt

